











## 6. References

1. Jain, L, Eaton DC. Physiology of fetal lung fluid clearance and the effect of labour. *Semin Perinatol* 2006; 30: 34-43.
2. Hansen AK, Wisborg K, Uldbjerg N, Henriksen TB. Elective caesarean section and respiratory morbidity in the term and near term neonate. *Acta Obstet Gynaecol Scan* 2007; 86: 389-94.
3. Kolas T, Saugstad OD, Daltveit AK, Nilsen ST, Oian P. Planned caesarean versus planned vaginal delivery at term: comparison of newborn infant outcomes. *Am J Obstet Gynecol* 2006; 195: 1538-43.
4. **O'Broovich H. Fluid clearance from the lungs of newborns, infants and children. *Paediatr Respir Rev* 2006; 7 (S1): S62-3.**
5. Doan E, Gibbons K, Tudehope D. The timing of elective caesarean deliveries and early neonatal outcomes in singleton infants born 37–41 weeks' gestation. *Aust N Z J Obstet Gynaecol*. 2014;54(4):340. [Available at: <https://doi.org/10.1111/ajo.12220>]
6. White SW, Newnham JP. Is it possible to safely prevent late preterm and early term births? *Semin Fetal Neonatal Med* 2019. [Available at: <https://pubmed.ncbi.nlm.nih.gov/30396760/>]
7. Bourke J, Wong K, Srinivasjois R, et al. Predicting Long-Term Survival Without Major Disability for Infants Born Preterm. *JAMA Pediatr* 2019;215:902019
8. Robinson m, Whitehouse AJO, Zubrick SR, Pennell CE, Jacoby P et al. *Aust N Z J Obstet Gynaecol*. 2013 Apr;53(2):143-51. [Available at: <https://pubmed.ncbi.nlm.nih.gov/23173808/>]
9. MacKay DF, Smith GCS, Dobbie R, Pell JP (2010). Delivery at 37 weeks' gestation is associated with a higher risk for child behavioural problems. *Aust N Z J Obstet Gynaecol*. 20rv-3.9871914(a)3.98277(l)356rv-367986.9











